



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

interesting results. All parts of *Coprinus* and other fungi exhibit distinct polarity, so that when parts of stems or pilei are grafted in their normal position on the corresponding parts of other individuals used as stocks, union readily takes place by anastomosing of the hyphae. When the part used as the scion is inserted in the reverse position, no union takes place. In partly resupinate forms, like *Polystictus*, polarity was exhibited in the same sense, but portions near the margins showed polarity to a less degree than the older parts.—H. HASSELBRING.

Lagenostoma.—This paleozoic type of seed, the first to be connected with Cycadofilicales, has been investigated further by Miss PRANKERD¹⁶ from preparations of *L. ovoides*. The *Lagenostoma* type of seed has peculiarities that are hard to relate to the structures of the more modern gymnosperm seeds, and any additional knowledge of the facts is welcome. The structures revealed by these new preparations are described in detail, and some interesting "theoretical suggestions" are made. The seed investigated strengthens the suggestion of OLIVER and SCOTT that the outer fleshy layer of the cycadean testa represents the cupule of *Lagenostoma*, the stony layer being developed after fusion of the *Lagenostoma* integument with the cupule. The method of pollination is discussed also, and the curious supposition that extraneous water must be brought to the pollen chamber for the swimming sperms is continued. The facts in reference to the peculiar "crevice-like" pollen chamber are somewhat cleared up. It is shown that the contact of the "central cone" with the outer layer of the nucellus is quite variable, so that apparent continuity might be developed in a variety of ways. The point of this is that a preparation showing a space below and continuity above does not prove necessarily that the pollen chamber is being formed from below upward. In certain specimens this very appearance was observed and yet there were pollen grains in the chamber. It is not even certain that the crevice-like chamber was continuous around the central cone. The specialized apical portion of the nucellus is called the "lagenostome," and the suggestion as to its morphology is very interesting. Miss PRANKERD sees in it a modified apical annulus, which in the fossil *Seftenbergia* is a multiseriate structure, but which in living forms has become simpler. If this be true, we have a fern connection for the structure that seemed to be hopelessly advanced, namely the seed.—J. M. C.

The foliar ray of dicotyledons.—BAILEY¹⁷ has followed up his previous work on the rays of certain groups by a more comprehensive study of the dicotyledons, resulting in some important conclusions. The primitive angio-

¹⁶ PRANKERD, THEODORA L., On the structure of the paleozoic seed *Lagenostoma ovoides* Will. Jour. Linn. Soc. London 40:461-490. pls. 22-24. figs. 3. 1912.

¹⁷ BAILEY, IRVING W., The evolutionary history of the foliar ray in the wood of the dicotyledons, and its phylogenetic significance. Ann. Botany 26:647-661. pls. 62, 63. 1912.